



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ALGEBRA.

$$81. \text{ Show that } \frac{a_1^r}{(a_1 - a_2)(a_1 - a_3)(a_1 - a_4) \dots (a_1 - a_n)} \\ + \frac{a_2^r}{(a_2 - a_1)(a_2 - a_3) \dots (a_2 - a_n)} + \dots + \frac{a_n^r}{(a_n - a_1)(a_n - a_2) \dots (a_n - a_{n-1})}$$

is zero if r is less than $n-1$; to 1 if $r=n-1$, and to $a_1 + a_2 + a_3 + \dots + a_n$ if $r=n$.

[*C. Smith's Treatise on Algebra.*]

$$82. \left. \begin{aligned} y^2 + yz + z^2 &= a^2 \\ z^2 + zx + x^2 &= b^2 \\ x^2 + xy + y^2 &= c^2 \end{aligned} \right\} \text{ find } x, y, \text{ and } z.$$

[*Ibid.*]

GEOMETRY.

83. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

θ being variable, find the locus of a point whose coördinates are

$$a \tan(\theta + \alpha), \quad b \tan(\theta + \beta).$$

84. Proposed by FREDERICK R. HONEY, Ph. B., New Haven, Conn.

Find the locus of a point which will trisect all arcs having a common chord.

85. Proposed by S. F. NORRIS, Professor of Astronomy and Mathematics, Baltimore City College, Baltimore, Md.

Prove by pure geometry. Give direct proof, if possible.

If the bisectors of two angles of a triangle are equal, the triangle is isosceles.

[From *Wentworth's Plane Geometry*, exercise 43, page 72.]

MECHANICS.

61. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

A body is suspended from a fixed point by an elastic string, which is stretched to double its natural length when the body is in equilibrium. Find how much the body must be depressed, so that when let go, it may just reach the point of suspension.

62. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

A particle of mass m moves in the circumference of an ellipse with constant rate v . It is constrained to move in that circumference by attractive forces in the two foci. To determine the magnitude of these forces.

DIOPHANTINE ANALYSIS.

58. Proposed by E. S. LOOMIS, Ph. D., Professor of Mathematics in Cleveland West High School, Berea, O.

"The base of a right-angled triangle is 105; find all the perpendiculars and hypotenuses to fit it, such that their values shall be integers."

59. Proposed by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics in Russell College, Lebanon, Va.

Find the sum of the m th powers of all the numbers less than P and prime to it, and then by substitution find the sum when $m=1, 2, 3, 4, 5$.